

Comparison of two Odonata communities from a natural and a modified rainforest in Papua New Guinea

Steffen Oppel

Wildlife Conservation Society, PO Box 277, Goroka, EHP, Papua New Guinea.
<steffen.oppel@gmx.net>

Present address: Department of Biology and Wildlife, 211 Irving 1, University of Alaska Fairbanks, Fairbanks, AK 99775, USA.

Key words: Odonata, dragonfly, species assemblage, forest modification, tropical rainforest, species richness and conservation, Papua New Guinea.

ABSTRACT

The Odonata fauna of Papua New Guinea is species-rich, but human population growth and resulting modification of primary rainforests may lead to a loss of valuable habitat and species diversity. In this study, I compared the odonate assemblages of a natural tropical rainforest and a modified forest in order to assess the loss that could result from human forest alteration. I collected odonates and recorded habitat use at both study sites for several weeks. The assemblages were compared with similarity indices, and total species richness was estimated using a jackknife procedure. The natural forest community, with 61 species, had both a higher diversity and evenness than the village community, with 38 species. Altogether I found 78 species of 13 families, of which 21 were shared between the two areas. Among the families with more than one species, Megapodagrionidae and Libellulidae had the highest similarity between the two sites, whereas Coenagrionidae and Platycnemididae had fairly dissimilar community composition. Three families occurred only in the natural forest. The most important habitats in the village were open sunny areas, artificial ditches, and small permanent creeks, compared to most running waters and the forest interior in the natural forest. Based on habitat preferences in the natural forest, species inhabiting temporary water sources under closed canopy rainforests are most vulnerable to forest modification. They comprised a third of the forest community, and I estimate that approximately 25% of natural rainforest species will disappear following human-induced forest alteration.

INTRODUCTION

Anthropogenic changes to tropical forest ecosystems have a large impact on native arthropod communities (Brown 1991). Deforestation is seen as a major threat to the conservation of odonate diversity in tropical forest landscapes worldwide (Clausnitzer 2004; Dijkstra & Vick 2004; Orr 2004; Paulson 2004; Rowe 2004).

In most cases, alteration or removal of the primary forest will lead to a reduction in species diversity and abundance, and communities in disturbed habitat will often be comprised of eurytopic and widespread generalists (Hill et al. 1995; Samways & Steytler 1996; Stewart & Samways 1998; Fermon et al. 2000; Willott et al. 2000; Clausnitzer 2003; Cleary et al. 2005). The influence of alterations on invertebrate communities is, however, difficult to predict. Disturbance may cause decreases or increases of species diversity in a certain area, as habitat modification may provide habitat for non-forest species (Owen 1971; Clausnitzer 2003; Ferreira-Peruquetti & Fonseca-Gessner 2003; Cleary et al. 2004; Hill & Hamer 2004). Furthermore, habitat modification may have different effects on species at different trophic levels (Petchey et al. 2004), and results of one taxon are generally imprecise predictors for effects on other taxa (Lawton et al. 1998). Studies on the impact of different forms of human forest modification on a variety of invertebrate groups are therefore urgently needed.

Papua New Guinea (PNG) is known for its very high biodiversity in many taxa, probably due to a rugged mountainous landscape that enhances local endemism (Lieftinck 1942, 1949; Heads 2001, 2002). The odonate fauna of PNG is poorly studied, and distribution and habitat requirements of many species are unknown (Brooks & Richards 1992; Polhemus 1995; Mack 1998; Rowe 2004). As in many other tropical countries the human population in Papua New Guinea is growing rapidly. This increases the pressure on the remaining natural areas and water sources (Henderson 1997). Human modification of the landscape usually results in deforestation and degradation of streams and rivers as a result of pollution and erosion, with negative consequences for stream-dwelling organisms (Timm et al. 2001; Buss et al. 2002).

In this study I compared the Odonata communities of a natural tropical rainforest and a nearby village. The village was characterised by subsistence gardening and small-scale forest clearing, and thus represented a typical modification of the forest landscape following human population growth and expansion. Besides comparing the communities as a whole and on the family level, I recorded habitat use of all species to determine which habitat types host assemblages most vulnerable to deforestation. The results are discussed to highlight habitats and species of the highest conservation concern.

MATERIAL AND METHODS

Study location

The two study sites were situated in the Crater Mountain Wildlife Management Area (6°43'S, 145°05'E), a 2,700 km² tract of continuous rainforest on the southern scarp of the central mountain range of Papua New Guinea. The climate at the two sites was stable throughout the year, and both rainfall and temperature did not exhibit seasonal patterns. The natural study site (Crater Mountain Biological Research Station – CMBRS) covered ca 2.5 km², ranging from 850 m to 1,300 m a.s.l., with aseasonal rainfall of ca 6.5 m per annum (Wright et al. 1997). The site

was entirely forested, except a small clearing for the research station and a helicopter landing pad. Water sources included one small, fish-free pond (0.25 ha), as well as innumerable streams and rivers with a high variability of flow rate, depending on rainfall. The forest floor was moist and muddy, and puddles formed quickly after rain and persisted for several days.

The village of Herowana was ca 10 km east of CMBRS, and consisted of a small grassy airstrip, subsistence gardens, small coffee plantations and scattered huts comprising an area of ca 4 km². Native primary and secondary forest was still present, but had been degraded by cutting. The village was ca. 1,300 m in elevation and featured several creeks and ditches, as well as a fish-free pond in semi-natural forest and a sedge swamp in abandoned gardens. The gardens were situated along a deep river valley, which descended to an elevation of 950 m a.s.l. Thus, the village site was on average at a slightly higher elevation. Open areas had mainly firm clay soil where puddles formed after rain and persisted for several days.

Sampling methods

I sampled odonates at CMBRS between November 2003 and August 2004, and in Herowana from July through August and in October 2004. Due to the aseasonal climate I assumed that the little temporal overlap of sampling periods did not significantly alter my results. The sampling period at CMBRS was about three times as long (112 days) as in Herowana (36 days). Due to a higher proportion of rare species the species richness of natural tropical forests is often underestimated at a given level of sampling effort (Morse et al. 1988; Godfray et al. 1999). Preliminary analysis suggested that the species accumulation curves of both sites showed a similar asymptotic tendency at the end of each sampling period (Opell 2005). Species accumulation curves are useful tools to standardize results from areas with unequal sampling effort (Moreno & Halffter 2000; Willott 2001; Ugland et al. 2003; Colwell et al. 2004), and I conclude that the different sampling effort does not have a significant effect on the results.

Sampling was carried out by slowly walking through all habitat types. Specimens were collected with butterfly nets and killed in jars with ethanol. They were stored in paper envelopes and kept in a box filled with silica gel and naphthalene. J. Michalski (Morristown, USA) and T.W. Donnelly (Binghampton, USA) examined the specimens, and a number of species still awaits formal identification or description.

I defined 10 distinct habitat types to record habitat use: open sunny areas without water (habitat type 1); forest interior without obvious surface water (2); temporary puddles in the forest interior (3); temporary puddles in open sunny areas (4); fish-free permanent pond (5); large, open and sunny river > 8 m wide (6), small, partially shaded river 4-8 m wide (7); small, permanent, mostly shaded creek 1-4 m wide (8); small, temporary creek in the forest < 1 m wide (9); and artificial ditch (10). The water in all running water sources (habitat types 6-9) was normally clear, except after heavy rain when waters were murky due to the sediment load. For every encounter with an adult odonate the habitat type and the number of individuals were recorded.

Analysis

To estimate total species richness I used the first-order jackknife estimator (Heltsh & Forrester 1983), as this was the most accurate and least biased estimator in a comparison of several procedures (Palmer 1990). I divided the species list of each area into shared species (shared with the other area) and only species (not shared with the other area), and calculated the percentage of shared and only species per habitat type. I then compared the communities of both sites using a new derivative of Jaccard's similarity index that is robust towards unequal sampling size (Chao et al. 2005). This index uses the relative abundance of all shared species, and I calculated similarity indices for all families and all habitat types using the total number of individuals per family or habitat type to calculate relative abundances. I assigned every species to a 'preferred' habitat type, even though a measure of habitat availability, required for true preference analysis (Jones 2001), was not feasible in this study. I defined 'preferred habitat' as the habitat type where the species was most commonly found, and omitted species with equal proportions of observations in different habitat types. Similarity indices, evenness, and species richness estimators were calculated using the program EstimateS (Colwell 2004), using the first-order jackknife estimate procedure.




Figure 1: Rank-abundance curves for Odonata species — ▲ in natural rainforest (CMBRS: Crater Mountain Biological Research Station), and ○ in modified rainforest (Herowana) in the Crater Mountain Wildlife Management Area, Papua New Guinea, in 2004. Note that for simplicity only the first 38 species are shown at CMBRS. See Table 2 for species identities.

Table 1. Inventory data of the two odonate communities — from a natural rainforest (CMBRS: Crater Mountain Biological Research Station), and a modified rainforest (Herowana) in the Crater Mountain Wildlife Management Area, Papua New Guinea, in 2004.

	CMBRS	Herowana
Number species	61	38
Number families	13	10
Anisoptera	16 (27%)	9 (24%)
Zygoptera	45 (73%)	29 (76%)
Species total		78
Shared species		21
Species locally endemic	40	17
Jaccard Abd. (Chao et al. 2005)		0.477
Simpson-Index	11.48	8.59
Evenness	0.804	0.715

RESULTS

Community comparison

I recorded a total of 78 species from 13 families in both study areas together, with 21 species occurring both in Herowana and at CMBRS (Table 1). A complete species list of the two areas is given in Table 2. The natural rainforest site had a higher number of species and families, and a higher evenness. A rank-abundance curve showed that the eight most common species contained a higher proportion of individuals in Herowana than at CMBRS (Fig. 1). Both areas had about three times as many Zygoptera species as Anisoptera species.

The estimated species richness calculated was 73 (± 3.55 s.d.) species at CMBRS and 53 (± 4.14 s.d.) species in Herowana. The observed species richness was therefore 84% and 72% of the estimated species richness at CMBRS and in Herowana, respectively. The estimator of the Jaccard abundance index including shared unseen species was 0.503.

For any given individual from either community, the chance that it belonged to a shared species was 0.477 (Jaccard Abundance-Index, Table 1). The similarity of the two communities was marginally different for Anisoptera (0.449) and for Zygoptera (0.499), but differed widely between families. Four families with one member each occurred in both study areas and had therefore a similarity of 1.0 (Calopterygidae, Chlorocyphidae, Protoneuridae, and Aeshnidae). Three families occurred only at CMBRS (Lestidae, Platystictidae, and Synthemistidae), and one family (Isostictidae) did not have a species common between the two study areas. Among the remaining families the Megapodagrionidae had a very high proportion of shared species individuals (0.724), followed by the Libellulidae (0.491). Coenagrionidae, Platycnemididae, and Corduliidae had dissimilar assemblages in the two study areas with similarities of 0.193, 0.144 and 0.145, respectively.

Table 2. List of all Odonata species — recorded in natural rainforest (CMBRS: Crater Mountain Biological Research Station), and in modified rainforest (Herowana) in the Crater Mountain Wildlife Management Area, Papua New Guinea, in 2004. For every species the preferred habitat type, number of observed individuals (no.) and the abundance rank are given. See text for the definition of preference. Species without preferred habitat type occurred in equal proportions in different habitat types. Note that abundances result from different sampling periods in the two study areas and might not be comparable in absolute terms.

Species name	CMBRS			Herowana		
	Preferred habitat	No.	Rank	Preferred habitat	No.	Rank
Calopterygidae						
<i>Neurobasis</i> sp. nov.	8	207	1	8	52	7
Chlorocyphidae						
<i>Rhinocypha tincta</i> Rambur	8	58	13	8	66	4
Lestidae						
<i>Indolestes tenuissimus</i> Tillyard	5	132	7		0	
Megapodagrionidae						
<i>Argiolestes kirbyi</i> Förster	7	78	10	8	76	3
<i>microstigma</i> Lieftinck	8	23	19		0	
<i>saltuarius</i> Lieftinck		0		8	16	9
<i>sidonia</i> Martin	8	40	16	8	14	10
sp. nov. A	7	9	28	10	3	19
sp. nov. B	9	3	39		0	
sp. C	9	2	45	8	6	15
sp. D	2	45	1	1	26	
<i>Podopteryx selysi</i> Förster	2	3	39		0	
Coenagrionidae						
<i>Palaiargia charmosyna</i> Lieftinck		0		2	2	21
<i>halcyon</i> Lieftinck		0		6	1	26
sp. A	8	3	39	8	1	26
sp. B	0		10	2	21	
<i>Papuagrion auriculatum</i> Lieftinck		0		1	1	26
<i>degeneratum</i> Lieftinck		0		8	1	26
<i>flavipedum</i> Lieftinck		0		10	1	26
<i>occipitale</i> Selys		0		8	1	26
sp. nov. A	2	6	35		0	
sp. B	2	10	25	1	1	26
sp. C	7	2	45		5	17
sp. D	8	32	10	2	21	
<i>Pseudagrion</i> sp. A		7	1	53		0
sp. B	9	1	53		0	
sp. nov. C		0		8	13	11
<i>Teinobasis dominula</i> Lieftinck		0		4	1	26
<i>scintillans</i> Lieftinck	9	43	15	10	90	2
sp. A	8	1	53		0	
sp. B	7	34		0		
sp. C	5	85	9		0	
sp. D	3	1	53		0	
<i>stigmatizans</i> Lieftinck		0		10	2	21
<i>Xiphiagrion cyanomelas</i> Selys	5	151	6		0	
<i>truncatum</i> Lieftinck		0		5	10	12

Species name	CMBRS			Herowana		
	Preferred habitat	No.	Rank	Preferred habitat	No.	Rank
Isostictidae						
<i>Selysionera cornelia</i> Lieftinck	7	17	21		0	
sp. A	1	53		0		
sp. nov. B		0		8	1	26
Platycnemididae						
<i>Arrhenicnemis</i> sp.	9	1	53		0	
<i>Idiocnemis australis</i> Gassmann	2	11	24		0	
<i>leonorae</i> Lieftinck		8	32		0	
<i>obliterata</i> Lieftinck	3	9	28		0	
<i>strumidens</i> Lieftinck	2	62	12		0	
<i>Paramecocnemis stilla-cruroris</i> Lieftinck	0		7	1	26	
<i>Rhyacocnemis prothoracica</i> Lieftinck	19	20	10	3	19	
<i>Torrenticnemis filicornis</i> Lieftinck		0			1	26
Gen. nov., sp. nov.	8	9	28		0	
Platystictidae						
<i>Drepanosticta</i> sp. A	2	12	23		0	
sp. B	7	2	45		0	
sp. C	7	4	38		0	
sp. D	6	3	39		0	
sp. E	9	5	37		0	
sp. F	7	1	53		0	
sp. G	9	1	53		0	
sp. H	9	9	28		0	
sp. I	7	2	45		0	
sp. J	8	2	45		0	
sp. K	2	1	53		0	
Protoneuridae						
<i>Nososticta finisterrae</i> Förster	7	190	5	8	44	8
Aeshnidae						
<i>Anax selysi</i> Förster	5	101	8	4	8	13
Corduliidae						
<i>Macromia</i> sp.	6	33	18	1	2	21
<i>Procordulia leopoldi</i> Fraser	5	194	2		0	
Synthemistidae						
<i>Synthemis primigenia</i> Förster	3	3	39		0	
Libellulidae						
<i>Agrionoptera longitudinalis</i> Selys	9	6	35		5	17
<i>Bironides glochidion</i> Lieftinck	8	3	39		0	
<i>Diplacina callirrhoe</i> Lieftinck	8	10	25	2	8	13
<i>smaragdina</i> Selys	6	36	17		0	
<i>Huonia hylophila</i> Lieftinck		0		10	59	6
<i>epinephela</i> Förster	6	194	3		0	
<i>Nannophlebia amphygyllis</i> Lieftinck	7	53	14		1	26
<i>Neurothemis stigmatizans</i> Fabricius	1	13	22		0	
<i>Orthetrum glaucum</i> Brauer	1	64	11	4	66	4
sp.	6	2	45		0	
<i>villosovittatum</i> Brauer	5	192	4	10	193	1
<i>Pantala flavescens</i> Fabricius		0		1	6	15
unidentified libellulid A	5	2	45		0	
unidentified libellulid B	5	10	25		0	

Habitat use comparison

In Herowana the most species-rich habitat types were ditches, small permanent creeks, and open sunny areas (Table 3). These three habitats also served as preferred habitat for 68% of all species (Fig. 2). At CMBRS, running water sources had the highest species richness, ranging from 18 species in small temporary creeks to 26 in small permanent creeks (Table 3). The distribution of preferred habitats was more balanced at CMBRS than in Herowana, with five habitat types being equally important (Fig. 2). These were the forest interior, the pond, small rivers, and small permanent and temporary creeks. They served as preferred habitat for 79% of all species.

The similarity of habitat specific odonate assemblages was generally smaller than the overall community similarity. In the forest interior and in sunny puddles, similarity was zero as no shared species occurred in those habitats in Herowana. Puddles in the forest and big rivers also had very low similarities of 0.090 and 0.145, respectively. The remaining habitat types had similarities ranging from 0.245 (pond) to 0.461 (small rivers).

In Herowana, more than half of the species preferring open sunny areas, the forest interior, sunny puddles, small permanent creeks and ditches also occurred at CMBRS. The remaining species in Herowana, showing a preference for the pond, large and small rivers did not occur at CMBRS. At CMBRS the distribution of shared and only species was different, and the highest ratio of shared species was found in small rivers (45%), small permanent creeks (50%), and in open sunny areas (50%). Only one (13%) and three (33%) of those species preferring the forest interior and small temporary creeks at CMBRS were shared with the site in Herowana.




Figure 2: Proportion of all Odonata species preferring a particular habitat type — □ in modified rainforest (Herowana), and ■ in natural rainforest (CMBRS: Crater Mountain Biological Research Station), Crater Mountain Wildlife Management Area, Papua New Guinea, in 2004. See text for definition of habitat types and preference.

Table 3. Number of species occurring in and preferring each habitat type — in a natural rainforest (CMBRS: Crater Mountain Biological Research Station), and in a modified rainforest (Herowana) in the Crater Mountain Wildlife Management Area, Papua New Guinea, in 2004. Shared species indicates the number of identical species occurring in a particular habitat type in both study areas. Preferred habitat is defined as the habitat type in which the most individuals of a species were found (see text for details).

Habitat type	Shared species	CMBRS		Herowana	
		Occurring	Preferring	Occurring	Preferring
Open sunny areas (1)	2	5	2	16	5
Forest interior (2)	0	11	8	6	2
Puddles in the forest interior (3)	1	7	2	8	0
Puddles in open sunny areas (4)	0	2	0	9	3
Pond (5)	2	8	8	4	1
Big river (6)	4	13	5	10	1
Small river (7)	3	21	11	5	1
Small permanent creek (8)	5	26	10	17	12
Small temporary creek (9)	3	18	9	7	0
Ditch (10)	0	0	0	21	9

In Herowana, the two most species-rich habitat types (small permanent creeks and ditches) hosted 53% of the only species, whereas at CMBRS five habitat types (forest interior, forest puddles, pond, small rivers, small temporary creeks) had fairly equal proportions of only species and a total of 63%. The forest interior had the largest proportion of only species (18% of all only species).

DISCUSSION

The odonate community in the natural rainforest at CMBRS is more diverse in species than in the modified forest and garden landscape in Herowana. Even though a longer sampling period in Herowana might have led to a higher number of recorded species, the estimated species richness suggests that the natural forest would still harbor substantially more species.

Overall species diversity of the two sites was high, especially when compared with studies of larger areas in Africa (Samways 1989; Clausnitzer 1999; Vick 1999, 2002; Dijkstra & Lempert 2003) and the Neotropics (Novelo Gutiérrez et al. 1988; de Marmels 1998; Ramírez et al. 2000; Paulson 2002). Species turnover between the sites was much lower than between the natural forest and another primary forest survey in the Lakekamu Basin in lowland PNG (Mack 1998), ca 200 km to the south-east. While decreasing similarity between distant sites is a common biogeographic pattern (Nekola & White 1999; Arita & Rodriguez 2002; Cleary & Genner 2004), it might also indicate that rates of local endemism in PNG could be equally high as in similar environments in South-East Asia (Orr 2003, 2004). This hypothesis might also be supported by the fact that species preferring

larger watercourses in Herowana were absent from structurally very similar rivers at CMBRS, although anisopterans are generally good dispersers that should be able to colonize areas only 10-15 km apart.

The dominance structure of the community in Herowana was more skewed than at CMBRS, resulting in a lower evenness score. This indicates that the modification of the rainforest results in an altered species assemblage, where a few generalist species benefit from large open areas and achieve dominance in the odonate community (Samways & Steytler 1996; Stewart & Samways 1998; Clausnitzer 2003). Many libellulids are known to be eurytopic generalists occurring over large geographic ranges (Silsby 2001). The family Libellulidae had a fairly high proportion of shared species between the two study sites, and the two most common species in Herowana (*Orithetrum villosovittatum* and *O. glaucum*) were together with *Agrionoptera longitudinalis* also present in the survey at Lakekamu Basin (Mack 1998). Coenagrionidae and Platynemididae had very low similarity scores. This might be a result of the high diversification within these families, which would confine most species to a very narrow ecological niche or geographic area. As a consequence, the ecological differences between CMBRS and Herowana will cause a relatively high species turnover within those families. The absence of the genus *Drepanosticta* from Herowana is, however, intriguing. While the species limits of this genus are currently unresolved (Orr 2004), the variety of habitats in which morphospecies of this genus were found suggests that the genus is adaptable to a broad spectrum of habitats.

Open sunny areas, artificial ditches, and temporary puddles in sunny areas were much more widespread and preferred by more species in Herowana than at CMBRS. In the natural forest the range of habitats preferred was wider and more evenly distributed, with species-rich assemblages found in running waters (rivers and creeks) as well as in the forest interior.

The species recorded in the forest interior away from obvious water sources might have been either foraging, or might depend on phytotelmata or water-filled tree-holes, as is known from other tropical regions (Lounibos et al. 1987; Kitching & Orr 1996; Louton et al. 1996; Clausnitzer 1997; Kitching 2000; Yanoviak 2001). *Podopteryx selysi* is known as a phytotelmata specialist (Kitching 2000), and this species was absent from Herowana. Most of the forest interior and small temporary water species at CMBRS did not occur in Herowana. Species depending on a closed canopy and temporary water sources are most susceptible to forest modification (Clausnitzer 2003), and the modification of the rainforest in Herowana might have led to drier conditions that do not meet the moisture requirements of certain species (Holdsworth & Uhl 1997; Silsby 2001; Cleary et al. 2004). Half of the species occupying the permanent creeks at CMBRS were also present in Herowana, indicating that these species are more tolerant to minor habitat modifications than species depending on temporary water sources.

Reduced shading of the water course may lead to an upward shift of certain species in disturbed landscapes (Korkeamäki & Suhonen 2002; Dijkstra & Lempert 2003). Species that are not dependent on a closed canopy may therefore tolerate habitat modification by moving upstream into shadier sections of the watercourse. Opening of the canopy along creeks may benefit species that are naturally limited by the availability of sunny areas along creeks. In Herowana, some creek species like *Rhinocypha tinctoria* were more abundant than in the natural forest, maybe as a result of extended creek sections being exposed to sunlight.

This study corroborates previous findings that deforestation and habitat disturbance lead to a lower β -diversity, and a species turnover towards more widespread generalists (Hill et al. 1995; Stewart & Samways 1998; Willott et al. 2000; Clausnitzer 2003; Samways 2003; Cleary et al. 2005). The higher dominance of generalist predators may furthermore cause a suppression of smaller species suffering from increased predation, especially in the larval stages (Cleary et al. 2004).

In this study, species associated with temporary puddles in the forest, small temporary creeks, and the forest interior comprised almost a third of the total odonate assemblage in the natural rainforest, and most of them could not be found in the modified forest. This suggests that modification of the forest caused by human subsistence gardening may lead to a loss of approximately 25% of the species associated with closed canopy forest habitats.

Given the potential for high levels of local endemism in PNG, forest modification might render many rainforest odonates vulnerable to extinction (Korkeamäki & Suhonen 2002). Conservation of these species will be difficult if only a few large nature reserves are set aside, which might be outside the natural range of many species. Conservation of odonates in PNG is therefore dependent on the preservation of natural rainforests in all parts of the country.

ACKNOWLEDGEMENTS

This study was initiated with the aid of the Wildlife Conservation Society, and was supported by the International Dragonfly Fund and the Worldwide Dragonfly Association. The Department of Environment and Conservation of Papua New Guinea gave generous permission to collect and export sample specimens of Odonata. Fieldwork was carried out with the permission of the Pawaian and Gimi people inhabiting the two study areas. I am very grateful to both groups for being supportive and hosting me on their land. I would like to thank Andy Mack and Ross Sinclair (WCS) for assistance and logistical support while doing the fieldwork. Stephen Richards and Bert Orr shared many valuable thoughts and thus improved the design and analysis of the study. Species identification would have been impossible without the help of John Michalski and Nick Donnelly, who processed the collected specimens, and I greatly appreciate their effort. The manuscript benefited from comments and suggestions by Bert Orr, Richard Rowe, Göran Sahlén, and Frank Suhling.

REFERENCES

- Arita, H.T. & P. Rodriguez, 2002. Geographic range, turnover rate and the scaling of species diversity. *Ecography* 25: 541-550.
- Brooks, S.J. & S.J. Richards, 1992. A new species of *Oreagrion* (Odonata: Coenagrionidae): montane damselflies from New Guinea. *Tijdschrift voor Entomologie* 135: 141-144.
- Brown, K.S., 1991. Conservation of neotropical environments: insects as indicators. In: Thomas, J.A. (ed.) "The conservation of insects and their habitats", Academic Press, London, pp. 349-404.

- Buss, D.F., D.F. Baptista, M.P. Silveira, J.L. Nessimian & L.F.M. Dorville, 2002. Influence of water chemistry and environmental degradation on macroinvertebrate assemblages in a river basin in south-east Brazil. *Hydrobiologia* 481: 125-136.
- Chao, A., R.L. Chazdon, R.K. Colwell & T.-J. Shen, 2005. A new statistical approach for assessing similarity of species composition with incidence and abundance data. *Ecology Letters* 8: 148-159.
- Clausnitzer, V., 1997. Reproductive behaviour of *Notiothemis robertsi* Fraser (Anisoptera: Libellulidae). *Odonatologica* 26: 451-457.
- Clausnitzer, V., 1999. Dragonfly (Odonata) records of Kakamega Forest, western Kenya, with notes on the ecology of rain forest species. *Journal of East African Natural History* 88: 17-24.
- Clausnitzer, V., 2003. Dragonfly communities in coastal habitats of Kenya: indication of biotope quality and the need of conservation measures. *Biodiversity and Conservation* 12: 333-356.
- Clausnitzer, V., 2004. Critical species of Odonata in eastern Africa. *International Journal of Odonatology* 7: 189-206.
- Cleary, D.F.R. & M.J. Genner, 2004. Changes in rain forest butterfly diversity following major ENSO-induced fires in Borneo. *Global Ecology & Biogeography* 13: 129-136.
- Cleary, D.F.R., A.Ø. Mooers, K.A.O. Eichhorn, J. van Tol, R. de Jong & S.B.J. Menken, 2004. Diversity and community composition of butterflies and odonates in an ENSO-induced fire affected habitat mosaic: a case study from East Kalimantan, Indonesia. *Oikos* 105: 426-446.
- Cleary, D.F.R., T.J.B. Boyle, T. Setyawati & S.B.J. Menken, 2005. The impact of logging on the abundance, species richness and community composition of butterfly guilds in Borneo. *Journal of Applied Entomology* 129: 52-59.
- Colwell, R.K., 2004. EstimateS: statistical estimation of species richness and shared species from samples. University of Connecticut, Storrs, CT. <<http://purl.oclc.org/estimates>>.
- Colwell, R.K., C.X. Mao & J. Chang, 2004. Interpolating, extrapolating, and comparing incidence-based species accumulation curves. *Ecology* 85: 2717-2727.
- de Marmels, J., 1998. A five year survey of an odonate community inhabiting a North Venezuelan mountain stream. *Odonatologica* 27: 189-199.
- Dijkstra, K.-D.B. & J. Lempert, 2003. Odonate assemblages of running waters in the Upper Guinean forest. *Archiv für Hydrobiologie* 157: 397-412.
- Dijkstra, K.-D.B. & G.S. Vick, 2004. Critical species of Odonata in western Africa. *International Journal of Odonatology* 7: 229-238.
- Fermon, H., M. Waltert, T.B. Larsen, U. Dall'Asta & M. Mühlenberg, 2000. Effects of forest management on diversity and abundance of fruit-feeding nymphalid butterflies in south-eastern Cote d'Ivoire. *Journal of Insect Conservation* 4: 173-188.
- Ferreira-Peruquetti, P.S. & A.A. Fonseca-Gessner, 2003. Comunidade de Odonata (Insecta) em áreas naturais de Cerrado e monocultura no nordeste do Estado de Sao Paulo, Brasil: relação entre o uso do solo e a riqueza faunística. *Revista Brasileira de Zoologia* 20: 219-224.
- Godfray, H.C.J., O.T. Lewis & J. Memmott, 1999. Studying insect diversity in the tropics. *Philosophical Transactions of the Royal Society of London (B)* 354: 1811-1824.
- Heads, M., 2001. Birds of paradise, biogeography and ecology in New Guinea: a review. *Journal of Biogeography* 28: 893-927.
- Heads, M., 2002. Birds of paradise, vicariance biogeography and terrane tectonics in New Guinea. *Journal of Biogeography* 29: 1-23.
- Heltsh, J.F. & N.E. Forrester, 1983. Estimating species richness using the jackknife procedure. *Biometrics* 39: 1-12.
- Henderson, M., 1997. Forest futures for Papua New Guinea: logging or community forestry? In: Clerk, C. (ed.) "Environment and development in the Pacific Islands", University of Papua New Guinea Press, Port Moresby.

- Hill, J.K. & K.C. Hamer, 2004. Determining impacts of habitat modification on diversity of tropical forest fauna: the importance of spatial scale. *Journal of Applied Ecology* 41: 744-754.
- Hill, J.K., K.C. Hamer, L.A. Lace & W.M.T. Banham, 1995. Effects of selective logging on tropical forest butterflies on Buru, Indonesia. *Journal of Applied Ecology* 32: 754-760.
- Holdsworth, A.R. & C. Uhl, 1997. Fire in Amazonian selectively logged forest and the potential for fire reduction. *Ecological Applications* 7: 713-725.
- Jones, J., 2001. Habitat selection in avian ecology: a critical review. *Auk* 118: 557-562.
- Kitching, R.L., 2000. Food webs and container habitats: the natural history and ecology of phytotelmata. Cambridge University Press, Cambridge, UK.
- Kitching, R.L. & A.G. Orr, 1996. The foodweb from water-filled treeholes in Kuala Belalong, Brunei. *Raffles Bulletin of Zoology* 44: 405-413.
- Korkeamäki, E. & J. Suhonen, 2002. Distribution and habitat specialization of species affect local extinction in dragonfly Odonata populations. *Ecography* 25: 459-465.
- Lawton, J.H., D.E. Bignell, B. Bolton, G.F. Bloemers, P. Eggleton, P.M. Hammond, M. Hodda, R.D. Holt, T.B. Larsen, N.A. Mawdsley, N.E. Stork, D.S. Srivastava & A.D. Watt, 1998. Biodiversity inventories, indicator taxa and effects of habitat modification in tropical forests. *Nature* 391: 72-75.
- Lieftinck, M.A., 1942. The dragonflies (Odonata) of New Guinea and neighbouring islands. Part VI. Results of the Third Archbold Expedition 1938-'39 and the Le Roux Expedition 1939 to Netherlands New Guinea (I. Anisoptera). *Treubia* 18: 441-608.
- Lieftinck, M.A., 1949. The dragonflies (Odonata) of New Guinea and neighbouring islands. Part VII. Results of the Third Archbold Expedition 1938-'39 and the Le Roux Expedition 1939 to Netherlands New Guinea (II. Zygoptera). *Nova Guinea (N.S.)* 5: 1-271.
- Lounibos, L.P., J.H. Frank, C.E. Machado Allison, P. Ocanto & J.C. Navarro, 1987. Survival, development and predatory effects of mosquito larvae in Venezuelan phytotelmata. *Journal of Tropical Ecology* 3: 221-242.
- Louton, J., J. Gelhaus & R. Bouchard, 1996. The aquatic macrofauna of water-filled bamboo (Poaceae: Bambusoideae: Guadua) internodes in a Peruvian lowland tropical forest. *Biotropica* 28: 228-242.
- Mack, A.L., 1998. A biological assessment of the Lakekamu Basin, Papua New Guinea. Conservation International, Washington, D.C.
- Moreno, C.E. & G. Halffter, 2000. Assessing the completeness of bat biodiversity inventories using species accumulation curves. *Journal of Applied Ecology* 37: 149-158.
- Morse, D.R., N.E. Stork & J.H. Lawton, 1988. Species number, species abundance and body length relationships of arboreal beetles in Bornean lowland rainforest trees. *Ecological Entomology* 13: 25-37.
- Nekola, J.C. & P.S. White, 1999. The distance decay of similarity in biogeography and ecology. *Journal of Biogeography* 26: 867-878.
- Novelo Gutiérrez, R., O. Canul Gonzalez & J. Camal Mex, 1988. The Odonata from the State of Quintana Roo, Mexico (Insecta: Odonata). *Folia Entomologica Mexicana*: 13-68.
- Oppel, S., 2005. Odonata in the Crater Mountain Wildlife Management Area, Papua New Guinea. IDP-Report 7: 1-28.
- Orr, A.G., 2003. Dragonflies of Borneo. Natural History Publications, Kota Kinabalu.
- Orr, A.G., 2004. Critical species of Odonata in Malaysia, Indonesia, Singapore and Brunei. *International Journal of Odonatology* 7: 371-384.
- Owen, D.F., 1971. Species diversity in butterflies in a tropical garden. *Biological Conservation* 3: 191-198.
- Palmer, M.W., 1990. The estimation of species richness by extrapolation. *Ecology* 71: 1195-1198.

- Paulson, D.R., 2002. Odonata records from Nayarit and Sinaloa, Mexico, with comments on natural history and biogeography. *Odonatologica* 31: 359-370.
- Paulson, D.R., 2004. Critical species of Odonata in the Neotropics. *International Journal of Odonatology* 7: 163-188.
- Petchev, O.L., A.L. Downing, G.G. Mittelbach, L. Persson, C.F. Steiner, P.H. Warren & G. Woodward, 2004. Species loss and the structure and functioning of multitrophic aquatic systems. *Oikos* 104: 467-478.
- Polhemus, D.A., 1995. A preliminary biodiversity survey of aquatic heteroptera and other aquatic insect groups in the Kikori River Basin, Papua New Guinea. In: WWF (ed.) "Field survey of biodiversity in the Kikori River Basin, Papua New Guinea", World Wildlife Fund, Washington, D.C., pp. 33-47.
- Ramírez, A., D.R. Paulson & C. Esquivel, 2000. Odonata of Costa Rica: diversity and checklist of species. *Revista de Biología Tropical* 48: 247-254.
- Rowe, R.J., 2004. Conservation of Odonata in the South Pacific and Australasia. *International Journal of Odonatology* 7: 139-147.
- Samways, M.J., 1989. Taxon turnover in Odonata across a 3000 m altitudinal gradient in Southern Africa. *Odonatologica* 18: 263-274.
- Samways, M.J., 2003. Threats to the tropical island dragonfly fauna (Odonata) of Mayotte, Comoro archipelago. *Biodiversity and Conservation* 12: 1785-1792.
- Samways, M.J. & N.S. Steytler, 1996. Dragonfly (Odonata) distribution patterns in urban and forest landscapes, and recommendations for riparian management. *Biological Conservation* 78: 279-288.
- Silsby, J., 2001. *Dragonflies of the world*. CSIRO, Collingwood.
- Stewart, D.A.B. & M.J. Samways, 1998. Conserving dragonfly (Odonata) assemblages relative to river dynamics in an African savanna game reserve. *Conservation Biology* 12: 683-692.
- Timm, H., M. Ivask & T. Möls, 2001. Response of macroinvertebrates and water quality to long-term decrease in organic pollution in some Estonian streams during 1990-1998. *Hydrobiologia* 464: 153-164.
- Ugland, K.I., J.S. Gray & K.E. Ellingsen, 2003. The species-accumulation curve and estimation of species richness. *Journal of Animal Ecology* 72: 888-897.
- Vick, G.S., 1999. A checklist of the dragonflies of the South West province of Cameroon with a description of a new species of the genus *Phyllogomphus* Selys, 1854. *Odonatologica* 28: 219-256.
- Vick, G.S., 2002. Preliminary biodiversity assessment of the odonate fauna of the Takamanda Forest Reserve, Cameroon. IDF-Report 4: 1-10.
- Willott, S.J., 2001. Species accumulation curves and the measure of sampling effort. *Journal of Applied Ecology* 38: 484-486.
- Willott, S.J., D.C. Lim, S.G. Compton & S.L. Sutton, 2000. Effects of selective logging on the butterflies of a Bornean rainforest. *Conservation Biology* 14: 1055-1065.
- Wright, D., H.J. Jessen, P. Burke & H.G. de Silva Garza, 1997. Tree and liana enumeration and diversity on a one-hectare plot in Papua New Guinea. *Biotropica* 29: 250-260.
- Yanoviak, S.P., 2001. Predation, resource availability, and community structure in Neotropical water-filled tree holes. *Oecologia* 126: 125-133.